

- ergic reticular system: Neocortical, subcortical and olfactory projections. *Brain*, 1967, 90, 497-520.
- Simpson, J. B. Subfornical organ involvement in angiotensin-induced drinking. In G. Peters, J. T. Fitzsimons, & L. Peters-Haefeli (Eds), *Control mechanisms of drinking*. Berlin: Springer-Verlag, 1975.
- Simpson, J. B. Localization of dipsogenic receptors for angiotensin II. In J. P. Buckley & C. M. Ferrario (Eds.), *Central actions of angiotensin and related hormones*. New York: Pergamon Press, 1977.
- Simpson, J. B., & Routtenberg, A. The subfornical organ and carbachol-induced drinking. *Brain Research*, 1972, 42, 135-157.
- Simpson, J. B., & Routtenberg, A. Subfornical organ: Site of drinking elicitation by angiotensin II. *Science*, 1973, 181, 1172-1175.
- Simpson, J. B., & Routtenberg, A. Subfornical organ: Acetylcholine elicits drinking. *Brain Research*, 1974, 79, 179-184.
- Simpson, J. B., & Routtenberg, A. Subfornical organ lesions reduce intravenous angiotensin-induced drinking. *Brain Research*, 1975, 88, 154-161.
- Simpson, J. B., Saad, W. A., & Epstein, A. N. The subfornical organ, the cerebrospinal fluid, and the dipsogenic action of angiotensin. In G. Onesti, K. E. Kim, & M. Fernandes (Eds), *Regulation of blood pressure by the central nervous system*. New York: Grune & Stratton, 1976.
- Sokabe, H., & Nakajima, T. Chemical structure and role of angiotensins in the vertebrates. *General and Comparative Endocrinology*, 1972, Supp. 3, 382-392.
- Spoerri, O. Über die Gefäßversorgung des Subfornikalorgans der Ratte. *Acta Anatomica*, 1963, 54, 333-348.
- Swanson, L. W., Marshall, G. R., Needleman, P., & Sharpe, L. G. Characterization of central angiotensin II receptors involved in the elicitation of drinking in the rat. *Brain Research*, 1973, 49, 441-446.
- Swanson, L. W., Sharpe, L. G., & Griffin, D. Drinking to intracerebral angiotensin II and carbachol: Dose-response relationships and ionic involvement. *Physiology and Behavior*, 1973, 10, 595-600.
- Tang, M., & Falk, J. Sar¹, ala⁸-angiotensin II blocks renin-angiotensin but not beta-adrenergic dipsogenesis. *Pharmacology Biochemistry and Behavior*, 1974, 2, 401-408.
- Thornborough, J. R., Passo, S. S., & Rothballe, A. B. Forebrain lesion blockade of the natriuretic response to elevated carotid blood sodium. *Brain Research*, 1973, 58, 355-363.

Received July 6, 1977 ■

Erratum to Wolgin and Teitelbaum

In the article "Role of Activation and Sensory Stimuli in Recovery from Lateral Hypothalamic Damage in the Cat" by David L. Wolgin and Philip Teitelbaum (*Journal of Comparative and Physiological Psychology*, 1978, Vol. 92, No. 3, pp. 474-500), three lines were printed incorrectly. On page 488, the last three sentences of the first paragraph in the right column should be two sentences and read as follows: The two cats with unilateral lesions displayed similar unilateral neglect, orienting to the mouse only in the ipsilateral visual field and preferentially using the ipsilateral forelimb to trap the mouse. Similar unilateral impairments have been reported in the rat (Marshall & Teitelbaum, 1974; Marshall et al., 1971).